ORIGINAL RESEARCH

Tree diversity drives diversity of arthropod herbivores, but successional stage mediates detritivores

Michael J. O'Brien^{1,2} Matteo Brezzi^{2,3} Andreas Schuldt⁴ Jia-Yong Zhang⁵ Keping Ma⁶ | Bernhard Schmid² | Pascal A. Niklaus²

¹Estación Experimental de Zonas Áridas, Consejo Superior de Investigaciones Científicas, Almería, Spain

²Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland

³Institute of Global Health, University of Geneva, Geneva, Switzerland

⁴German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

⁵Institute of Ecology, Zhejiang Normal University, Jinhua, Zhejiang Province, China ⁶State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing, China

Correspondence

Michael J. O'Brien, Estación Experimental de Zonas Áridas, Consejo Superior de Investigaciones Científicas, Almería, Spain. Email: mikey.j.obrien@gmail.com and

Pascal A. Niklaus, Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland. Email: pascal.niklaus@ieu.uzh.ch

Funding information

Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung, Grant/ Award Number: P2ZHP3_161986 and P300PA_167758; Universität Zürich, Grant/ Award Number: University Research Priority Program on Global Change and Biodiversity; Deutsche Forschungsgemeinschaft, Grant/ Award Number: DFG FOR 891 TP1; European project BACCARA funded by the European Commission's Seventh Framework Program (FP7/2007-2013), under Grant Agreement No. 226299; National Science Foundation of China grant (NSFC 30710103907 and 30930005)

Abstract

The high tree diversity of subtropical forests is linked to the biodiversity of other trophic levels. Disentangling the effects of tree species richness and composition, forest age, and stand structure on higher trophic levels in a forest landscape is important for understanding the factors that promote biodiversity and ecosystem functioning. Using a plot network spanning gradients of tree diversity and secondary succession in subtropical forest, we tested the effects of tree community characteristics (species richness and composition) and forest succession (stand age) on arthropod community characteristics (morphotype diversity, abundance and composition) of four arthropod functional groups. We posit that these gradients differentially affect the arthropod functional groups, which mediates the diversity, composition, and abundance of arthropods in subtropical forests. We found that herbivore richness was positively related to tree species richness. Furthermore, the composition of herbivore communities was associated with tree species composition. In contrast, detritivore richness and composition was associated with stand age instead of tree diversity. Predator and pollinator richness and abundance were not strongly related to either gradient, although positive trends with tree species richness were found for predators. The weaker effect of tree diversity on predators suggests a cascading diversity effect from trees to herbivores to predators. Our results suggest that arthropod diversity in a subtropical forest reflects the net outcome of complex interactions among variables associated with tree diversity and stand age. Despite this complexity, there are clear linkages between the overall richness and composition of tree and arthropod communities, in particular herbivores, demonstrating that these trophic levels directly impact each other.

KEYWORDS

BEF-China, biodiversity, canopy layers, community composition, ecosystem functioning, forest succession, plant-herbivore interactions, trophic groups

1 | INTRODUCTION

et al., 2006; Hooper et al., 2005; Isbell et al., 2015; Siemann, Tilman, Haarstad, & Ritchie, 1998). Associations between the diversity of plants and other trophic levels have been studied intensively in grasslands (Haddad et al., 2009; Scherber et al., 2010; Siemann et al., 1998),

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2017 The Authors. Ecology and Evolution published by John Wiley & Sons Ltd.

Plant diversity is important for maintaining ecosystem functioning

and for supporting the diversity of other trophic levels (Balvanera

but the relationship between plant and animal diversity in forests has received far less attention. Most research examining the effect of plant diversity on animal communities in forests has focused on the loss of tree species due to land-use change and logging (Dunn, 2004; Edwards et al., 2010; Ewers et al., 2015; Fredericksen & Fredericksen, 2004; Tews et al., 2004). However, understanding the variables that drive the distribution, diversity and abundance of trophic levels that depend upon forest vegetation in a contiguous forest landscape can provide insights into the factors that promote biodiversity and ecosystem functioning in subtropical forests.

An important relationship in terrestrial ecosystems is that between plants and arthropods because of the feedbacks that exist between these groups of organisms. Plants provide habitat and food while arthropods may alter plant diversity (Bagchi et al., 2014; Have et al., 2006; Kempel et al., 2015), contribute to decomposition (Donoso, Johnston, Clay, & Kaspari, 2013), mediate plant reproduction (Gonzalez-Megias, 2016; Zvereva, Lanta, & Kozlov, 2010), and disperse seeds (Kalisz, Hanzawa, Tonsor, Thiede, & Voigt, 1999). Relationships between plant and arthropod diversity may depend on the dominant plant life form in the ecosystem—i.e., forests dominated by trees and grasslands dominated by herbaceous plants may have different plant– arthropod interactions.

Forests have physical attributes for arthropod communities that are different from grasslands because of their structural complexity, which may supersede the effects of plant diversity on arthropod diversity (Southwood, Brown, & Reader, 1979). Forests have high spatial heterogeneity with horizontal variation (gap dynamics) in stem densities, light and temperature (Chazdon & Fetcher, 1984; Chen et al., 1999; Raich, 1989), and vertical variation in guality and guantity of leaf and woody tissues (Ellsworth & Reich, 1993). Vertical canopy strata can have direct effects on arthropod communities. For example, birds and other predators in the canopy may reduce abundances of pollinators and herbivores (Van Bael, Brawn, & Robinson, 2003) and variation in light and microclimate can change the nutrients in leaf tissues (Le Corff & Marguis, 1999). Variation in abiotic conditions such as soil characteristics and climatic variables associated with altitude and topography (Paoli, 2006; Paoli, Curran, & Zak, 2006; Pendry & Proctor, 1997; Proctor, Lee, Langley, Munro, & Nelson, 1988) may also influence arthropod communities independent of tree diversity. In addition, trees produce recalcitrant tissues that decompose slowly and create biotope space that persists for long times. For example, fallen or standing dead plant material provides space for breeding and larval development regardless of the surrounding living tree species (Irmler, Heller, & Warning, 1996; Jacobs, Spence, & Langor, 2007; Schiegg, 2000). Combined, these variables make forests distinct from grasslands, which may alter plant-arthropod relationships.

Furthermore, the long-lived nature of trees means that a forest consists of a mosaic of stand age classes due to disturbances that occur at different spatial and temporal scales (Bergeron, 2000). Large canopy gaps promote the recruitment of early-successional tree species with traits for establishment and rapid growth in high light environments while undisturbed areas will have long-lived species with traits that support shade tolerance and stress resistance (lida et al., 2014; Kohyama, Suzuki, Partomihardjo, Yamada, & Kubo, 2003). Environmental conditions in older forest stands will have decreased light and soil drying and understories will be cooler than younger forest stands in gaps (Chen et al., 1999; Raich, 1989). Therefore, stand age, which correlates with functional traits, environmental conditions, and the quantity of woody debris (Chen et al., 1999; Jacobs et al., 2007; Raich, 1989), may mediate arthropod diversity across the land-scape more than tree species diversity.

These factors of stand age, heterogeneity in spatial structure and environmental conditions, may supersede the effects of tree diversity on arthropod diversity in forests. However, the strength and magnitude of the tree-arthropod diversity relationship may also differ between different arthropod functional groups occupying different trophic niches. For example, detritivores will respond to food quantity and quality that may be less related to living tree diversity than to species identity or functional traits of dead plant material that effect detritus quantity and quality (Donoso et al., 2013; Graça, Pozo, Canhoto, & Elosegi, 2002; Hansen, 2000; Hättenschwiler & Jørgensen, 2010). In contrast, herbivore richness and abundance should be more directly linked to living tree diversity because herbivores feed on these plants (Andow, 1991; Knops et al., 1999). Predators, in turn, may be indirectly linked to plants through alterations in the diversity and abundance of herbivores (Hutchinson, 1959; Knops et al., 1999) and may therefore show a weaker treearthropod diversity relationship (Balvanera et al., 2006). Pollinators may operate independently of stand-level tree diversity altogether due to their dependence on flowering and potentially longdistance travel (Bawa, Bullock, Perry, Coville, & Grayum, 1985; Sobek, Tscharntke, Scherber, Schiele, & Steffan-Dewenter, 2009). Therefore, to determine the factors that mediate arthropod distributions across a forested landscape, arthropod functional groups must be assessed separately while accounting for tree diversity and forest age.

In this study, we tested the effect of forest variables (e.g., tree species diversity, stand age, and vertical position) and arthropod community characteristics (e.g., taxon richness and abundance) using a comparative study design across gradients of tree diversity and stand age in a subtropical forest. We selected forest plots so that these gradients were relatively independent, allowing us to separate the effects of tree diversity and stand age (Baruffol et al., 2013; Bruelheide et al., 2011). We sampled arthropods of four functional groups (i.e., detritivores, herbivores, pollinators, and predators) in the understorey and the canopy to assess differences in vertical position as well. We posit that these variables will differentially affect the different arthropod functional groups. We further assessed whether arthropod community compositions were related to tree community composition. Given the host specificity of some arthropods, we hypothesized that the species compositions of tree and arthropod communities should show different associations depending on the arthropod functional group considered (as outlined above).

2 | MATERIALS AND METHODS

2.1 | Experimental design

In June 2010, we placed arthropod traps in 27 plots of a comparative study in the Gutianshan National Nature Reserve (GNNR) in Zhejiang Province, China (Baruffol et al., 2013; Bruelheide et al., 2011; Castro-Izaguirre et al., 2016). The plots in the GNNR were established in 2008 to encompass a gradient of stand ages (20-120 years) and tree species richness levels (25-69 species of all trees greater than 10 cm diameter at breast height). Four to seven plots were assigned to five strata based on successional stage of the forests stands (<20 years; <40 years; <60 years; <80 years; >80 years). Within each stratum, plots spanned a gradient of lower to higher tree species diversity (Table S1). Bruelheide et al. (2011) presented the age and diversity spread of these plots, indicating their continuous distribution (see Figure 4 in Bruelheide et al. (2011) see Table S1). The forest is continuous (see Fig. S1) and classified as subtropical with ~2,000 mm of rain mainly falling from March to September (Yu, Hu, Yu, Ding, & Fang, 2001). The 27 plots are 30 × 30 m in size, and their altitude ranges from 250 to 900 m above sea level. The average distance among plots was 3.400 m (95% CI: 535-7420).

Plots were subdivided into nine 10 × 10 m subplots. In July and August 2010, ten yellow sticky traps (9 × 11 cm sticky area, double face, MIOPLANT, Switzerland) were placed in each study plot. Five traps were suspended by bamboo sticks, in the understorey, 2 m above ground with one trap placed in the middle subplot and one in each of the four corner subplots (at least 5 m from the plot edge). The sticky sides of the understorey traps faced east-west. Five additional traps were placed in the canopy, suspended from trees. One tree from each of the five most abundant species in the plot was chosen at random to suspend a trap in the canopy. Tree traps were positioned inside the middle of the crown of each tree between 3 and 18 m above the ground (tree heights ranged from 7 to 30 m) hanging from 0.35 mm fishing line and stabilized by small ballasts. Due to logistic constraints, trap exposure varied from 6 to 9 days depending on the plot. After collection, the traps were covered with plastic and stored in a freezer until further processing. Because yellow sticky traps present a sampling bias, absolute values of arthropod abundance cannot be estimated by our study. However, their use across all plots allows comparison of abundances among plots and should not inhibit the total number of species captured (Hoback, Svatos, Spomer, & Higley, 1999; Missa et al., 2009).

2.2 | Arthropod sorting and counting

Arthropods were identified directly on the traps and were classified by order and to morpho-species based on external morphological characteristics (Yuan, Zhang, Feng, & Hua, 2006; Zheng & Gui, 1999). Larvae were considered as separate morpho-species because their diet often differs from their respective adult form and the difficulty in defining larvae to the correct adult morpho-species. However, the ambiguity in larvae identification did not affect our ____Ecology and Evolution

results as larvae only represented 0.4% of the arthropods captured (123 individuals). Based on the inspection of the arthropod mouthparts, taxonomic experience, and known arthropod populations in the province, each morpho-species was assigned to one of six functional groups: (1) herbivore folivores, (2) herbivore sapsuckers, (3) predators (including parasitoids), (4) detritivores, (5) pollinators (nonherbivorous), and (6) miscellaneous arthropods that could not be classified more precisely. The use of mouth parts was employed for insects that could not be assigned to a family containing only one functional type. These initial groups were aggregated into four classes for analysis: (1) herbivores (folivores + sapsuckers), (2) predators, (3) pollinators, and (4) detritivores. Insects with ambiguous classification were set to miscellaneous and are not included in our analysis (17% of the total). In total, we collected 28,198 arthropods belonging to 17 different orders and 598 morphospecies (Table S2 and S3). Some morpho-species may have been wrongly assigned to a functional group because these assignments were based on taxonomy and morphology (of mouth parts), and direct observations of feeding behavior were not made (Table S4). Therefore, sensitivity analysis was performed whereby the orders Lepidoptera, Hemiptera, and Coleoptera were either removed or all designated as herbivores. This analysis showed that the results were not dependent upon the feeding group assignment of these orders (Table S5).

2.3 | Data analysis

We performed two sets of analyses: The first tested the effect of tree richness and stand age on arthropod richness and abundance, and the second assessed evidence for relationships between tree and arthropod compositions.

In order to assess the importance of tree diversity and stand age on arthropod diversity and abundance, we modeled arthropod richness and abundance, separately for each functional group, as a function of stand age (a continuous variable), tree species richness (a continuous variable, trees with a diameter at breast height of at least 10 cm), vertical position (a fixed factor with two levels; understorey and canopy), and all two-way interactions between the continuous variables and vertical position using a linear mixed-effects model. We also included a covariate for altitude. Because of low arthropod abundance on some traps, we aggregated traps from the same vertical position within plots which combined to 54 observations (2 sampling heights × 27 plots). To meet assumptions of linearity, arthropod richness and abundance were log-transformed after adding one to account for zeros. Plot (a factor with 27 levels) and trap exposure time (a factor with 22 levels) were fit as random terms.

To test for specific associations between arthropod and tree composition, we first calculated the dissimilarities in tree community composition among plots using the Jaccard index based on tree species presence-absence per plot (Fig. S2). We performed the same analysis on the arthropod community composition among plots for each functional group separately. These dissimilarity matrices were transformed into two-dimensional space by principal coordinate analysis (PCoA), and the first two axes were used as orthogonal metrics of differences in tree or arthropod community composition among plots. The Bray– Curtis index (with square-root transformed abundances) was also calculated for arthropod functional groups, but results were similar to those using the Jaccard index (Fig. S3–S6). Therefore, only results from analysis of the Jaccard data are discussed.

We performed constrained analysis of proximities on the Jaccard distance matrix of the arthropod community for each functional group to test the effect of stand age and tree composition on arthropod community composition. The PCoA ordination of the communities of each functional group was constrained by the two PCoA axes derived from the PCoA of tree composition, stand age, and tree species richness. We tested the significance of the constraining terms with a permutation test. If the inertia in the permuted models was lower than in the constrained model, then the association modeled was considered statistically significant.

We further tested the importance of tree composition for arthropod composition of each functional group using the PCoA axis 1 of tree composition as a predictor in a linear model with type-I sum of squares of each PCoA axis of the arthropod functional group compositions. Therefore, if tree composition affected the arthropod community composition of a functional group, then the PCoA axis of tree composition would significantly correlate with the first or second PCoA axis of the arthropod functional group. In other words, plots with a more similar tree community composition would have more similar arthropod community composition. Furthermore, we tested for the independence of tree composition from stand age by adding stand age in before tree composition, and if tree composition was independent of stand age, then its importance would be retained even after fitting stand age first in the model.

All linear and mixed models were performed with the asreml-R package (ASReml 3, VSN International, Hemel Hempstead, UK), using R 3.3.2 (http://r-project.org). The vegdist function in vegan package

(Oksanen et al., 2015) was used to calculate Jaccard dissimilarities. Principal coordinate ordination was carried out with the cmdscale function, and the constrained analysis of proximities was performed with the capscale function (Legendre & Anderson, 1999) in the vegan package.

3 | RESULTS

Arthropod richness and abundance were differentially affected by tree species richness, stand age, and vertical position, depending on the arthropod functional group (Figures 1 and 2; Tables S7 and S8). Herbivore richness significantly increased with tree species richness (Figure 1a) while herbivore abundance was significantly higher in the canopy than in the understorey (Figure 2a). In addition, stand age had a significant negative effect on herbivore richness despite a tendency for later successional stages to have higher tree species richness. Predator richness and abundance were significantly higher in the understorey than in the canopy (Figures 1b and 2b) and increased nonsignificantly with tree species richness. Detritivore richness and abundance significantly decreased with stand age and were significantly higher in the understory than in the canopy (Figures 1c and 2c). Detritivores were unaffected by tree species richness. In addition, detritivore abundance decreased significantly faster in the understorey than in the canopy as stand age increased (Figure 2c). Pollinator richness and abundance were not affected by any variables and were generally found in equal numbers in all plots. ANOVA tables for all richness and abundance analyses are in Tables S7 and S8.

The constrained analysis of proximities on the different arthropod communities indicated that only compositions of herbivore and detritivore groups were explained by tree species composition (Figs S3–S6), as evidenced by the PCoA axes of tree composition explaining herbivore composition (p < .1) and detritivore composition (p < .1). In



FIGURE 1 The effect of tree species richness and stand age on arthropod richness of different functional groups. (a) The relationship between herbivore richness and tree species richness for both understorey (black lines and points) and canopy (gray lines and points) traps. (b) The relationship between predator richness and tree species richness for understorey and canopy traps. (c) The relationship between detritivore richness and stand age in the understory and canopy traps. Arthropod richness was log-transformed but presented with axes back-transformed.

200

100

50

0

(a)

Herbivore abundance per plot



FIGURE 2 The effect of tree species richness and stand age on arthropod abundance of different functional groups. (a) Herbivore abundance increased with species richness in the canopy (gray lines and points) and understorey (black lines and points). (b) Predator abundance increased in the understorey and canopy traps but was always higher in the understorey. (c) Detritivore abundance declined with stand age but faster in understorey traps. Arthropod abundance was log-transformed but presented with axes back-transformed.

Tree species richness

our direct comparison of PCoA axis of tree composition to the PCoA axes of each arthropod functional group (Figs S7-S10), only herbivore composition was related to tree composition. The PCoA axis 2 of herbivore community composition was related to the tree community composition ($F_{1,25}$ = 4.4, *p* < .05 and $F_{1,25}$ = 2.5, *p* = .1 for tree composition fit before and after stand age, respectively; Figure 3). However, the community compositions of the other arthropod functional groups were not related to tree composition regardless of the inclusion of stand age in the model.

Tree species richness



FIGURE 3 Tree composition correlated with herbivore community composition. A relationship between principal coordinates of tree and herbivore community composition was found. Plots with more similar tree communities had more similar herbivore communities.

| DISCUSSION 4

In this assessment of arthropod communities across a gradient of tree species richness and stand age, we found that different arthropod functional groups showed different associations with the forest characteristics. Specifically, herbivore richness and abundance increased with tree species richness, and detritivore richness and abundance decreased with forest age. In contrast, the richness and abundance of predators and pollinators were not strongly associated with tree diversity or stand age. Therefore, higher tree diversity only sustained a higher diversity of herbivores, but stand age and vertical position superseded the effects of tree diversity on detritivores. Furthermore, a direct link between herbivore community composition and tree community composition was found, which suggests a potential role for species-specific interactions between herbivores and trees in this subtropical forest.

Stand age (years)

4.1 | Tree and herbivore diversity

Herbivores showed a clear relationship with tree species richness. Most likely the connection between species-rich tree communities and more diverse herbivore communities was due to more tree species supporting a greater array of feeding demands combined with the benefits provided by feeding on a diversity of plants that improves overall diet and fitness (Coley & Barone, 1996). Recent work by Brezzi, Schmid, Niklaus, and Schuldt (2017) showed higher levels of feeding on locally rare species, indicating that generalists or at least nonspecialists had a strategy to increase their diversity of food intake. Although only marginally significant, herbivore abundance also increased with tree diversity, in support of growing evidence that herbivory increases with tree species richness (Brezzi et al., 2017; Scherber et al., 2006; Schuldt et al., 2010; Vehviläinen, Koricheva, & Ruohomäki, 2007).

Herbivore community composition was also linked to tree community composition. Whether these results support the role of herbivores in promoting a diversity-productivity relationship depends on the feeding preferences of these herbivores (Barone, 1998). If host-specific feeding dominates or if host-specific feeders concentrate in low-diversity forest stands and generalists are more common in high-diversity stands, then herbivores would inhibit productivity of low-diversity forests and promote a biodiversity effect (Barone, 1998; Root, 1973; Schuldt et al., 2010; Zhang et al., 2017). Furthermore, recent research suggests that more diverse forest stands may have increased nutrient cycling rates due to faster leaf turnover, a pattern that may be mediated by higher herbivore diversity (Huang et al., 2017). Although our results cannot determine feeding preferences or underlying biodiversity mechanisms, they clearly show greater tree diversity sustains a more diverse herbivore community.

4.2 | Stand age and detritivore associations

Detritivore richness and abundance were negatively related to stand age. Early-successional forest stands have canopies dominated by fast growing light-demanding trees with slow-growing, shade-tolerant species recruiting underneath. Fast growing trees have higher leaf nutrient content and less recalcitrant foliage (Eichenberg, Trogisch, Huang, He, & Bruelheide, 2013; Garnier et al., 2004; Li, Pei, Kéry, Niklaus, & Schmid, 2017), and these characteristics promote higher quality detritus which would support greater diversity and abundance of detritivores (Cortez, Garnier, Pérez-Harguindeguy, Debussche, & Gillon, 2007). In addition, the understorey supported greater richness and abundance, which is likely due to the greater quantity of dead and down tissue than the canopy. The lack of evidence for a relationship between tree species diversity and detritivores and the marginal evidence for a role of tree composition supports our initial hypotheses that stand age can supersede the importance of tree species diversity.

4.3 | What shapes predator and pollinators communities?

Predator richness and abundance also tended to increase with tree species richness, although the relationship was statistically not significant. The weak response of predators to tree species richness may indicate an indirect effect mediated by the herbivore community (Balvanera et al., 2006; Knops et al., 1999). The pattern of herbivores and predators tracking the next lower trophic level (i.e., predators follow herbivores follow plants) supports the diversity-trophic structure hypothesis (Hutchinson, 1959; Knops et al., 1999; Murdoch, Evans, & Peterson, 1972). The weakening of the positive trend of predators with trees suggests a bottom-up effect in support of Balvanera et al. (2006). A positive signal with only 27 plots suggests that biologically tree diversity is showing important cascading effects on arthropod functional groups.

Pollinators in general were operating independent of any forest characteristics. We suspect pollinator beta-diversity is driven not by tree species diversity, but instead specifically by flowering diversity and phenology. Therefore, without data on the diversity of flowering species, we are unable to determine mechanisms promoting this functional group.

5 | CONCLUSION

In the subtropical forest studied here, more diverse tree communities hosted more diverse and abundant herbivore communities (and to a weaker extent predator communities), a pattern which was not found in other arthropod guilds. Tree and herbivore community compositions were also correlated, indicating a degree of specialized herbivory. Therefore, tree diversity drives herbivore diversity in spite of variables associated with stand age and spatial heterogeneity. In contrast, detritivores were responding to stand age which likely represents a proxy for functional traits associated with different tree growth strategies. Combined, these results indicate the importance of maintaining tree species and functional diversity across forest successional stages for promote arthropod diversity in subtropical forests.

ACKNOWLEDGMENTS

MOB was supported in part by the Stiefel-Zangger foundation and a Swiss National Science Foundation Post.Doc Mobility Fellowship (P2ZHP3_161986, P300PA_167758). MB was funded by a European project BACCARA from the European Commission's Seventh Framework Program (FP7/2007-2013) to BS (Grant Agreement No. 226299). AS, BS and PAN gratefully acknowledge funding by the German Research Foundation (DFG FOR 891 TP1). We further are indebted to Helge Bruelheide for his continuous efforts in running this research platform and acknowledge Gutianshan National Nature Reserve for support of the study. Thank you to two anonymous reviewers for their detailed comments on earlier versions. This research was supported by the University Research Priority Program on Global Change and Biodiversity at the University of Zurich.

CONFLICT OF INTEREST

None declared.

AUTHOR CONTRIBUTIONS

MOB analyzed the data, wrote the manuscript, and led the revisions. MB conceived and carried out the experiment and contributed to revisions. AS provided conceptual development with expertise on arthropods and contributed to revisions. JYZ contributed to experimental design and insect identification. KM provided logistical support for working in China and the setup of the larger experiment. BS and PAN wrote the grant that funded the project and contributed to the experimental design, analysis, writing, and revisions.

ORCID

Michael J. O'Brien D http://orcid.org/0000-0003-0943-8423

Ecology and Evolution

Bernhard Schmid D http://orcid.org/0000-0002-8430-3214

REFERENCES

- Andow, D. A. (1991). Vegetational diversity and arthropod population response. Annual Review of Entomology, 36, 561–586.
- Bagchi, R., Gallery, R. E., Gripenberg, S., Gurr, S. J., Narayan, L., Addis, C. E., ... Lewis, O. T. (2014). Pathogens and insect herbivores drive rainforest plant diversity and composition. *Nature*, 506, 85–88.
- Balvanera, P., Pfisterer, A. B., Buchmann, N., He, J. S., Nakashizuka, T., Raffaelli, D., & Schmid, B. (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecology Letters*, 9, 1146–1156.
- Barone, J. A. (1998). Host-specificity of folivorous insects in a moist tropical forest. *Journal of Animal Ecology*, *67*, 400–409.
- Baruffol, M., Schmid, B., Bruelheide, H., Chi, X., Hector, A., Ma, K., ... Niklaus, P. A. (2013). Biodiversity promotes tree growth during succession in subtropical forest. *PLoS One*, *8*, e81246.
- Bawa, K. S., Bullock, S. H., Perry, D. R., Coville, R. E., & Grayum, M. H. (1985). Reproductive biology of tropical lowland rain forest trees. II. Pollination systems. *American Journal of Botany*, 72, 346–356.
- Bergeron, Y. (2000). Species and stand dynamics in the mixed woods of Quebec's southern boreal forest. *Ecology*, *81*, 1500–1516.
- Brezzi, M., Schmid, B., Niklaus, P., & Schuldt, A. (2017). Tree diversity increases levels of herbivory damage in a subtropical forest canopy: Evidence for dietary mixing by arthropods? *Journal of Plant Ecology*, 10, 13–27.
- Bruelheide, H., Böhnke, M., Both, S., Fang, T., Assmann, T., Baruffol, M., ... Schmid, B. (2011). Community assembly during secondary forest succession in a Chinese subtropical forest. *Ecological Monographs*, 81, 25–41.
- Castro-Izaguirre, N., Chi, X., Baruffol, M., Tang, Z., Ma, K., Schmid, B., & Niklaus, P. A. (2016). Tree diversity enhances stand carbon storage but not leaf area in a subtropical forest. *PLoS One*, *11*, e0167771.
- Chazdon, R. L., & Fetcher, N. (1984). Photosynthetic light environments in a Lowland tropical rain forest in Costa Rica. *Journal of Ecology*, 72, 553–564.
- Chen, J., Saunders, S. C., Crow, T. R., Naiman, R. J., Brosofske, K. D., Mroz, G. D., ... Franklin, J. F. (1999). Microclimate in forest ecosystem and landscape ecology. *BioScience*, 49, 288–297.
- Coley, P. D., & Barone, J. A. (1996). Herbivory and plant defenses in tropical forests. Annual Review of Ecology and Systematics, 27, 305–335.
- Cortez, J., Garnier, E., Pérez-Harguindeguy, N., Debussche, M., & Gillon, D. (2007). Plant traits, litter quality and decomposition in a Mediterranean old-field succession. *Plant and Soil*, 296, 19–34.
- Donoso, D. A., Johnston, M. K., Clay, N. A., & Kaspari, M. E. (2013). Trees as templates for trophic structure of tropical litter arthropod fauna. *Soil Biology and Biochemistry*, 61, 45–51.
- Dunn, R. R. (2004). Managing the tropical landscape: A comparison of the effects of logging and forest conversion to agriculture on ants, birds, and lepidoptera. *Forest Ecology and Management*, 191, 215–224.
- Edwards, D. P., Hodgson, J. A., Hamer, K. C., Mitchell, S. L., Ahmad, A. H., Cornell, S. J., & Wilcove, D. S. (2010). Wildlife-friendly oil palm plantations fail to protect biodiversity effectively. *Conservation Letters*, *3*, 236–242.
- Eichenberg, D., Trogisch, S., Huang, Y., He, J. S., & Bruelheide, H. (2013). Shifts in community leaf functional traits are related to litter decomposition along a secondary forest succession series in subtropical China. *Journal of Plant Ecology*, 8, 401–410.
- Ellsworth, D. S., & Reich, P. B. (1993). Canopy structure and vertical patterns of photosynthesis and related leaf traits in a deciduous forest. *Oecologia*, *96*, 169–178.
- Ewers, R. M., Boyle, M. J. W., Gleave, R. A., Plowman, N. S., Benedick, S., Bernard, H., ... Turner, E. C. (2015). Logging cuts the functional importance of invertebrates in tropical rainforest. *Nature Communications*, *6*, 6836.

- Fredericksen, N. J., & Fredericksen, T. S. (2004). Impacts of selective logging on amphibians in a Bolivian tropical humid forest. *Forest Ecology and Management*, 191, 275–282.
- Garnier, E., Cortez, J., Billes, G., Navas, M.-L., Roumet, C., Debussche, M., ... Toussaint, J.-P. (2004). Plant functional markers capture ecosystem properties during secondary succession. *Ecology*, 85, 2630–2637.
- Gonzalez-Megias, A. (2016). Within- and trans-generational effects of herbivores and detritivores on plant performance and reproduction. *Journal of Animal Ecology*, 85, 283–290.
- Graça, M. A. S., Pozo, J., Canhoto, C., & Elosegi, A. (2002). Effects of Eucalyptus plantations on detritus, decomposers, and detritivores in streams. *The Scientific World*, 2, 1173–1185.
- Haddad, N. M., Crutsinger, G. M., Gross, K., Haarstad, J., Knops, J. M. H., & Tilman, D. (2009). Plant species loss decreases arthropod diversity and shifts trophic structure. *Ecology Letters*, 12, 1029–1039.
- Hansen, R. A. (2000). Effects of habitat complexity and composition on a diverse litter microarthropod assemblage. America, 81, 1120–1132.
- Hättenschwiler, S., & Jørgensen, H. B. (2010). Carbon quality rather than stoichiometry controls litter decomposition in a tropical rain forest. *Journal of Ecology*, 98, 754–763.
- Have, P., Effects, L., Ecosystem, O. N., By, P., Diversity, C. P., & Biomass, N. O. T. P. (2006). Predators have large effects on ecosystem properties by changing plant diversity, not plant biomass. *Control*, 87, 1432–1437.
- Hoback, W. W., Svatos, T. M., Spomer, S. M., & Higley, L. G. (1999). Trap color and placement affects estimates of insect family-level abundance and diversity in a Nebraska salt marsh. *Entomologia Experimentalis et Applicata*, 91, 393–402.
- Hooper, D. U., Chapin, F. III, Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., ... Wardle, D. A. (2005). Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, 75, 3–35.
- Huang, Y., Ma, Y., Zhao, K., Niklaus, P. A., Schmid, B., & He, J.-S. (2017). Positive effects of biodiversity on litterfall quantity and quality along a chronosequence in a subtropical forest. *Journal of Plant Ecology*, 10, 28–35.
- Hutchinson, G. E. (1959). Homage to Santa Rosalia or Why are there so many kinds of animals? *The American Naturalist*, *93*, 145–159.
- Iida, Y., Poorter, L., Sterck, F., Kassim, A. R., Potts, M. D., Kubo, T., & Kohyama, T. S. (2014). Linking size-dependent growth and mortality with architectural traits across 145 co-occurring tropical tree species. *Ecology*, 95, 353–363.
- Irmler, U., Heller, K., & Warning, J. (1996). Age and tree species as factors influencing the populations of insects living in dead wood (Coleoptera, Diptera: Sciaridae, Mycetophilidae). *Pedobiologia*, 148, 134–148.
- Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., ... Eisenhauer, N. (2015). Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature*, 526, 574–577.
- Jacobs, J. M., Spence, J. R., & Langor, D. W. (2007). Influence of boreal forest succession and dead wood qualities on saproxylic beetles. *Agricultural and Forest Entomology*, 9, 3–16.
- Kalisz, S., Hanzawa, F. M., Tonsor, S. J., Thiede, D. A., & Voigt, S. (1999). Ant-mediated seed dispersal alters spatial pattern of seed relatedness in an outcrossing population of Trillium grandiflorum. *Ecology*, 80, 2620–2634.
- Kempel, A., Razanajatovo, M., Stein, C., Unsicker, S. B., Auge, H., Weisser, W. W., ... Prati, D. (2015). Herbivore preference drives plant community composition. *Ecology*, 96, 2923–2934.
- Knops, J. M. H., Tilman, D., Haddad, N. M., Naeem, S., Mitchell, C. E., Haarstad, J., ... Groth, J. (1999). Effects of plant species richness on invasion dynamics, disease outbreaks, insect abundances and diversity. *Ecology Letters*, 2, 286–293.
- Kohyama, T., Suzuki, E., Partomihardjo, T., Yamada, T., & Kubo, T. (2003). Tree species differentiation in growth, recruitment and allometry in relation to height in a Bornean mixed dipterocarp forest. *Journal of Ecology*, 91, 797–806.

- Le Corff, J., & Marquis, R. J. (1999). Differences between understorey and canopy in herbivore community composition and leaf quality for two oak species in Missouri. *Ecological Entomology*, 24, 46–58.
- Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: Testing multispecies responses in multifactorial experiments. *Ecological Monographs*, 69, 1–24.
- Li, X., Pei, K., Kéry, M., Niklaus, P. A., & Schmid, B. (2017). Decomposing functional trait associations in a Chinese subtropical forest. *PLoS ONE*, 12, e0175727.
- Missa, O., Basset, Y., Alonso, A., Miller, S. E., Curletti, G., De Meyer, M., ... Wagner, T. (2009). Monitoring arthropods in a tropical landscape: Relative effects of sampling methods and habitat types on trap catches. *Journal of Insect Conservation*, 13, 103–118.
- Murdoch, W. W., Evans, F. C., & Peterson, C. H. (1972). Diversity and Pattern in Plants and Insects. *Ecology*, 53, 819–829.
- Oksanen, J., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R., ... Wagner, H. (2015) Vegan: Community ecology package.
- Paoli, G. D. (2006). Divergent leaf traits among congeneric tropical trees with contrasting habitat associations on Borneo. *Journal of Tropical Ecology*, 22, 397.
- Paoli, G. D., Curran, L. M., & Zak, D. R. (2006). Soil nutrients and beta diversity in the Bornean Dipterocarpaceae: Evidence for niche partitioning by tropical rain forest trees. *Journal of Ecology*, 94, 157–170.
- Pendry, C. A., & Proctor, J. (1997). Altitudinal zonation of rain forest on Bukit Belalong, Brunei: Soils, forest structure and floristics. *Journal of Tropical Ecology*, 13, 221–241.
- Proctor, J., Lee, Y. F., Langley, A. M., Munro, W. R. C., & Nelson, T. (1988). Ecological studies on Gunung Silam, a small ultrabasic mountain in Sabah, Malaysia. I. environment, forest structure and floristics. *Journal* of Ecology, 76, 320–340.
- Raich, J. W. (1989). Seasonal and spatial variation in the light environment in a tropical dipterocarp forest and gaps. *Biotropica*, 21, 299–302.
- Root, R. B. (1973). Organization of a plant-arthropod association in simple and diverse habitats: The fauna of Collards (Brassica Oleracea). *Ecological Monographs*, 43, 95–124.
- Scherber, C., Eisenhauer, N., Weisser, W. W., Schmid, B., Voigt, W., Fischer, M., ... Tscharntke, T. (2010). Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment. *Nature*, 468, 553–556.
- Scherber, C., Mwangi, P. N., Temperton, V. M., Roscher, C., Schumacher, J., Schmid, B., & Weisser, W. W. (2006). Effects of plant diversity on invertebrate herbivory in experimental grassland. *Oecologia*, 147, 489–500.
- Schiegg, K. (2000). Effects of dead wood volume and connectivity on saproxylic insect species diversity. *Ecoscience*, 7, 290–298.
- Schuldt, A., Baruffol, M., Böhnke, M., Bruelheide, H., Härdtle, W., Lang, A. C., ... Assmann, T. (2010). Tree diversity promotes insect herbivory in subtropical forests of south-east China. *Journal of Ecology*, 98, 917–926.

- Siemann, E., Tilman, D., Haarstad, J., & Ritchie, M. (1998). Experimental tests of the dependence of arthropod diversity on plant diversity. *The American Naturalist*, 152, 738–750.
- Sobek, S., Tscharntke, T., Scherber, C., Schiele, S., & Steffan-Dewenter, I. (2009). Canopy vs. understory: Does tree diversity affect bee and wasp communities and their natural enemies across forest strata? *Forest Ecology and Management*, 258, 609–615.
- Southwood, T., Brown, V., & Reader, P. (1979). The relationships of plant and insect diversities in succession. *Biological Journal of the Linnean Society*, 12, 327–348.
- Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M. C., Schwager, M., & Jeltsch, F. (2004). Animal species diversity driven by habitat heterogeneity/diversity: The importance of keystone structures. *Journal of Biogeography*, 31, 79–92.
- Van Bael, S. A., Brawn, J. D., & Robinson, S. K. (2003). Birds defend trees from herbivores in a Neotropical forest canopy. *Proceedings of the National Academy of Sciences*, 100, 8304–8307.
- Vehviläinen, H., Koricheva, J., & Ruohomäki, K. (2007). Tree species diversity influences herbivore abundance and damage: Meta-analysis of long-term forest experiments. *Oecologia*, 152, 287–298.
- Yu, M.-J., Hu, Z.-H., Yu, J.-P., Ding, B.-Y., & Fang, T. (2001). Forest vegetation in Gutianshan Natural Reserve in Zhejiang. *Journal of Zhejiang University*, 27, 375–380.
- Yuan, F., Zhang, Y., Feng, J., & Hua, B. (2006). *Taxonomy of Hexapoda*, 2nd ed. Beijing: China Agricultural University Press.
- Zhang, J., Bruelheide, H., Chen, X., Eichenberg, D., Kröber, W., Xu, X., ... Schuldt, A. (2017). Tree diversity promotes generalist herbivore community patterns in a young subtropical forest experiment. *Oecologia*, 183, 455–467.
- Zheng, L., & Gui, H. (1999). *Insect Classification, Part I & II*, 1st ed. Nanjing: Nanjing Normal University Press.
- Zvereva, E. L., Lanta, V., & Kozlov, M. V. (2010). Effects of sap-feeding insect herbivores on growth and reproduction of woody plants: A metaanalysis of experimental studies. *Oecologia*, 163, 949–960.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: O'Brien MJ, Brezzi M, Schuldt A, et al. Tree diversity drives diversity of arthropod herbivores, but successional stage mediates detritivores. *Ecol Evol*. 2017;00:1–8. https://doi.org/10.1002/ece3.3411